

New Data to the Knowledge on the Corticicolous Mite Fauna in Hungary (Acari: Prostigmata, Astigmata, Oribatida)

G. RIPKA¹, A. FAIN² and H. R. BOLLAND³

¹Plant Health and Soil Conservation Station of Budapest, H-1519 Budapest, Hungary

²Institut royal des Sciences naturelles de Belgique, B-1000 Bruxelles, Belgium

³Institute of Systematics and Population Biology, University of Amsterdam,
1098 SM Amsterdam, The Netherlands

Authors give a report on the results of the regular collections made between 1990 and 1998 on ornamental trees and shrubs, on streets, in parks, in green spaces of housing estates, in private gardens and in arboreta. A total of 19 corticolous mite species were found belonging to 14 families. *Michaelopus corticalis* (Michael, 1885), *Hericia hericia* Robin, 1868, *Eupalopsis maseriensis* (Canestrini et Fanzago, 1876), *Mediolata vandergeesti* (Gomaa et Bolland, 1982), *Cunaxoldes kielczewskii* Michocka, 1982, and *Neophyllobius* spec. nov. are reported for the first time for the Hungarian fauna. *Hemisarcopeltis budensis* Fain et Ripka, 1998, *M. corticalis* and *Cheletogenes ornatus* (Canestrini et Fanzago, 1876) were the most frequent species in the acarofauna associated with scale insects.

Key words: Corticolous Mites, Acari, Prostigmata, Astigmata, Oribatida

Among arboreal mites corticolous mites are less known contrary to foliage-dwelling mites. In the last three decades, Lebrun (1976), André (1986, 1987), Momen (1987, 1988), Momen and Lundqvist (1996), Suski (1972), Doberski (1986), Bolland (1983, 1986, 1991), Smiley (1992), Kehl and Weigmann (1992), Weigmann and Jung (1992) studied the mites on bark of trees and shrubs. Their results proved the considerable richness of corticolous acarofauna.

A part of cortex-dwelling mites is algae and lichens feeder. An other group is fungivorous and saprophagous. The third group is predaceous upon the smallest arthropods. In Belgium, André (1986) reported 22 species of Tydeidae and 6 species of Stigmaeidae from bark of different tree species. In Bulgaria, Nachev and Trenchev (1987), while in Italy, Vacante (1989) as well as Vacante and Gerson (1987) reported the mites found associated with coccids from fruit trees. Nachev and Trenchev (1987) observed *Thyreophagus entomophagus* (Laboulbene), *Cheletogenes ornatus* (Canestrini et Fanzago), *Cheyletia flabellifera* (Michael), *Hemisarcopeltis malus* (Shimer), *Amblyseius cucumeris* (Oudemans) and *Bdella iconica* Berlese under the shields of *Sphaerolecanium prunastri* (Fonscolombe), *Parlatoria oleae* (Colvée), *Quadraspidiotus marani* Zahradník and *Quadraspidiotus pyri* (Lichtenstein). *Th. entomophagus* was the most frequent species. Vacante (1989) reported *H. malus* from *Lepidosaphes beckii* (Newman), *Aonidiella aurantii* (Maskell), *Parlatoria pergandei* Comstock and *Aspidiotus nerii* Bouché. Several members of Stigmaeidae, Eupalopsellidae, Cunaxidae and Camerobiidae are predators of armored scale insects and can be found on the bark of trees (Cooreman,

1958, Meyer and Ueckermann, 1984, Bolland, 1986, Vacante and Gerson, 1987). Smiley and Knutson (1983), also Gerson (1973, 1985) summarized the knowledge on the role and biology of predatory mites of coccids. Fain et al. (1995) as well as Fain and Ripka (1998) reported new species of the genus *Hemisarcopeltis* that predaceous upon diaspidids.

The present knowledge of corticolous mites in Hungary remains scanty. Balogh and Mahunka (1980) reported oribatids, Komlovszky (1980, 1984) recorded cunaxid and cheyletid species. Lucza et al. (1996) found 4 cheyletid species on tree bark infested with armored scale insects. Ripka et al. (1997) collected the following 6 tarsonemid species: *Tarsonemus nodosus* Schaarschmidt, 1959, *Tarsonemus lobosus* Suski, 1965, *Tarsonemus karli* Sharonov et Mitrofanov, 1986, *Tarsonemus waitei* Banks, 1912, *Dendroptus willmanni* (Schaarschmidt, 1959) and an other *Dendroptus* sp. together with coccids. *Aceria heteronyx* (Nalepa, 1891) causes small stem galls on twigs and branches of *Acer platanoides*. An other eriophyoid mite, *Acalitus phloeocoptes* (Nalepa, 1890) causes stem galls on *Prunus domestica* and *Prunus spinosa* (Ripka and de Lillo, 1997). Ripka and Kaźmierski (1998a) recorded the following 8 tydeid species in association with scale insects: *Lorryia reticulata* (Oudemans, 1928), *Lorryia obliqua* (Kuznetzov, 1973), *Lorryia* cf. *woolleyi*, *Triophtydeus immanis* Kuznetzov, 1973, *Triophtydeus triophthalmus* (Oudemans, 1929), *Tydeus caudatus* (Duges, 1834), *Tydeus californicus* (Banks, 1904) and *Homeopronematus staerkii* (Schrufft, 1972). Out of them *T. immanis* was the most frequent mite. The following two tydeids: *Lorryia reticulata* and *L. obliqua* were found solely on bark. Ripka and Kaźmierski (1998b) found *Mediolata mariaefrancae* André, 1977 on the bark of tree infested with scale insects.

Materials and Methods

Between 1990 and 1998, a mite survey was made on ornamental trees and shrubs of streets, squares, parks, botanical gardens, private gardens, and green areas of housing estates in all districts of Budapest (in some other localities in Hungary and Croatia, also herbaceous plants were occasionally examined). Plant samples from 301 woody plant species (minimum 5 shoots, branches or bark/plant or 25 leaves/plant) were taken in plastic bags from January to late November. During plant examination with binocular microscope (upper and lower surfaces of the leaves, petioles, buds, bark, flowers, galls, etc.), all the mites found on the plant samples were put into AGA solution or directly into lactic acid. After the clearing of mites in the lactic acid, they were placed into Keifer's medium 2 following the Keifer's mounting method for eriophyoid mites (Keifer, 1952). The cleaned and dyed mites were then placed into Keifer's medium 3. The Hoyer's medium with sorbitol (Keifer, 1975), and Heinze's polyvinyl alcohol (PVA) medium (Schmutterer, 1959) were used alternatively for mounting the specimens. The slide preparations were dried in thermostat at 32 °C and then sealed with nail varnish. Specimens were examined with a phase contrast microscope. For mite determination the keys of Fain (1982), Bolland (1991), Smiley (1992) as well as several original species descriptions and other available reports were used.

Results

Corticolous mites were collected on 48 woody plant species out of 301 studied belonging to 19 families. These mites were found in 90 plant samples. A total of 19 mite species were identified belonging to 14 families (see Table 1 for the exact host/locality data). The following species were new for the Hungarian fauna: *Cunaxoides kielczewskii* Michocka, *Eupalopsis maseriensis* (Canestrini et Fanzago), *Hericia hericia* Robin, *Mediolata vandergeesti* (Gomaa et Bolland), *Michaelopus corticalis* (Michael). *M. corticalis*, *Hemisarcoptes budensis* Fain et Ripka and *Cheletogenes ornatus* were the most frequent species in the mite fauna associated with scale insects. Corticolous mites occurred on 16 species of family Rosaceae, on 8 species of family Oleaceae, and on 4 species of family Salicaceae. *Ch. ornatus* was collected on coccid infested cortex, exceptionally on leaves. *H. budensis*, *C. kielczewskii*, *E. maseriensis*, *M. mariae francae*, *M. vandergeesti*, *Neophyllobius* sp. near *sturmerwoodi* and *Neophyllobius* spec. nov. were found on branches infested with the following scale insects: *Pseudaulacaspis pentagona* (Targioni-Tozzetti), *Unaspis euonymi* (Comstock), *Epidiaspis leperii* (Signoret), *Chionaspis salicis* (Linnaeus), *Aulacaspis rosae* (Bouché), *Diaspidiotus gigas* (Thiem et Gerneck), *Diaspidiotus ostreaeformis* (Curtis), *Diaspidiotus perniciosus* (Comstock), and *Lepidosaphes ulmi* (Linnaeus) (for more data on Coccoidea see Ripka et al., 1996). The cortex-dwelling camerobiid, cheyletid, cunaxid, eupalopsellid and stigmaeid mites usually occurred in low numbers, but the coccid infestations were severe. Phoretic hypopodes (entomophilous deuteronymphs) of *H. budensis* were collected on *Chilocorus renipustulatus* (Scriba) and *Chilocorus bipustulatus* (L.) (Coleoptera: Coccinellidae).

M. corticalis was found under the scales of dead *P. pentagona*, *E. leperii*, *U. euonymi*, *Ch. salicis*, *Diaspidiotus marani* Zahradník, *D. gigas*, *D. perniciosus*, *L. ulmi* and colonies of *Pseudochermes fraxini* (Kaltenbach). *H. hericia* was collected under the peeling (loose) bark of London plane trunk. Since the paper of Ripka et al. (1997) an other tarsonemid mite was found associated with coccid. *Tarsonemus bachmaieri* Cooreman, 1958 was found under the scales of *Carulaspis juniperi* (Bouché). Oribatids were found only on bark of twigs in parks, cemetery and arboreta. They could not be found on street trees. The oribatid mites do not tolerate the aerial pollution in urban habitat. The vast majority of corticolous mite species is slow-moving, exception *T. immanis*.

Table 1

Corticolous mite species collected from ornamental trees and shrubs
(* = new species for the Hungarian fauna)

Mite species	Plant species	Sampling place	Sampling date
Acaria: Prostigmata			
fam. Cheyletidae			
<i>Cheletacarus raptor</i> Volgin, 1961			
	<i>Populus x canescens</i>	Óbuda	25.08. 1998
	<i>Populus simonii</i>	Vérmező	25.01. 1991
	<i>Rhamnus catharticus</i>	Józsefváros	04.01. 1995
<i>Cheletogenes ornatus</i> (Canestrini et Fanzago, 1876)			
	<i>Celastrus orbiculatus</i>	Gellérthegy	09.09. 1997
	<i>Crataegus laevigata</i>	Gellérthegy	05.02. 1991
	<i>Ilex aquifolium</i>	Pestszentlőrinc	02.09. 1992
	<i>Jasminum beesianum</i>	Városliget	04.02. 1995
	<i>Populus x canescens</i>	Óbuda	25.08. 1998
	<i>Populus simonii</i>	Sasad	13.09. 1991
	<i>Populus tremula</i>	Újpalota	30.08. 1994
	<i>Prunus avium</i>	Kispest	02.04. 1992
	<i>Thuja occidentalis</i>	Budaörs (Pest c.)	07.03. 1995
<i>Prosocheyla traubi</i> (Baker, 1949)			
	<i>Populus simonii</i>	Sasad	13.09. 1991
fam. Eupalopsellidae			
* <i>Eupalopsis maseriensis</i> (Canestrini et Fanzago, 1876)			
	<i>Crataegus laevigata</i>	Gellérthegy	24.01. 1995
	<i>Crataegus monogyna</i>	Gellérthegy	26.01. 1995
	<i>Malus pumila</i> var. <i>niedzwetzkyana</i>	Gellérthegy	26.01. 1995
fam. Stigmaeidae			
<i>Mediolata mariae-francae</i> André, 1977			
	<i>Euonymus europaeus</i>	Gellérthegy	15.06. 1997
	<i>Jasminum beesianum</i>	Gellérthegy	27.01. 1995
	<i>Rhamnus catharticus</i>	Józsefváros	04.01. 1995
	<i>Tilia x euchlora</i>	Józsefváros	29.06. 1994
* <i>Mediolata vandergeesti</i> (Gomaa et Bolland, 1982)			
	<i>Crataegus monogyna</i>	Gellérthegy	26.01. 1995
	<i>Ribes uva-crispa</i>	Széchenyi-hegy	04.02. 1991
fam. Camerobiidae			
* <i>Neophyllobius</i> sp. near <i>sturmerwoodi</i> Bolland, 1991			
	<i>Juglans nigra</i>	Kőbánya	29.03. 1995
* <i>Neophyllobius</i> sp. nov.			
	<i>Euonymus europaeus</i>	Gellérthegy	15.06. 1997
	<i>Euonymus europaeus</i>	Gellérthegy	09.09. 1997

Table 1 (cont.)

Mite species	Plant species	Sampling place	Sampling date
fam. Cunaxidae			
* <i>Cunaxoides kielczewskii</i> Michocka, 1982			
	<i>Picea pungens</i>	Orosháza (Békés c.)	07.07. 1998
	<i>Rhus typhina</i>	Kőbánya	04.01. 1995
	<i>Taxus baccata</i>	Orbán-hegy	25.08. 1994
fam. Tarsonemidae			
<i>Tarsonemus bachmaieri</i> Cooreman, 1958	<i>Thuja</i> sp.	Józsefváros	28.10. 1991
<i>Tarsonemus nodosus</i> Schaarschmidt, 1959	<i>Celastrus orbiculatus</i>	Gellérthegy	09.04. 1997
	<i>Euonymus europaeus</i>	Rákoskeresztúr	17.08. 1997
		Rákoskeresztúr	17.10. 1997
<i>Tarsonemus waitei</i> Banks, 1912	<i>Euonymus europaeus</i>	Rákoskeresztúr	17.08. 1997
		Rákoskeresztúr	17.10. 1997
Acari: Astigmata			
fam. Hemisarcopidae			
<i>Hemisarcopes budensis</i> Fain et Ripka, 1998			
	<i>Celastrus orbiculatus</i>	Gellérthegy	09.09. 1997
	<i>Crataegus laevigata</i>	Gellérthegy	24.01. 1995
	<i>Euonymus europaeus</i>	Gellérthegy	27.01. 1995
	<i>Fraxinus angustifolia</i>	Gellérthegy	15.06. 1997
		Víziváros	08.02. 1991
		Óbuda	11.09. 1991
	<i>Fraxinus pennsylvanica</i>	Víziváros	08.02. 1991
	<i>Malus baccata</i>	Gellérthegy	24.01. 1995
	<i>Phellodendron amurense</i>	Margitsziget	21.02. 1991
	<i>Populus x canescens</i>	Óbuda	03.04. 1991
		Óbuda	25.08.1998
	<i>Populus simonii</i>	Lágymányos	05.02. 1991
		Sasad	13.09. 1991
	<i>Prunus domestica</i>	Újpalota	12.02. 1991
	<i>Pyrus betulifolia</i>	Gellérthegy	18.01. 1995
	<i>Pyrus pyraster</i>	Józsefváros	04.01. 1995
	<i>Rhus typhina</i>	Kőbánya	04.01. 1995
	<i>Sophora japonica</i>	Víziváros	11.02. 1991
fam. Acaridae			
* <i>Michaelopus corticalis</i> (Michael, 1885)			
	<i>Acer platanoides</i>	Orosháza (Békés c.)	07.07. 1998
	<i>Acer pseudoplatanus</i>	Csongrád (Csongrád c.)	26.05. 1998
	<i>Celastrus orbiculatus</i>	Gellérthegy	07.08. 1998
	<i>Celastrus scandens</i>	Józsefváros	04.01. 1995
	<i>Cornus alba</i>	Órmező	28.03. 1995

Table 1 (cont.)

Mite species	Plant species	Sampling place	Sampling date
	<i>Cornus stolonifera</i>	Órméző	10.02. 1995
	<i>Crataegus laevigata</i>	Gellérthegy	24.01. 1995
	<i>Crataegus monogyna</i>	Gellérthegy	26.01. 1995
	<i>Crataegus succulenta</i>	Gellérthegy	24.01. 1995
	<i>Daphne mezereum</i>	Gellérthegy	24.01. 1995
	<i>Euonymus europaeus</i>	Gellérthegy	27.01. 1995
	<i>Forsythia suspensa</i>	Pasarét	09.09. 1997
		Belváros	17.09. 1990
	<i>Fraxinus excelsior</i>	Pasarét	10.04. 1993
	<i>Ilex aquifolium</i>	Városliget	17.05. 1991
	<i>Jasminum beesianum</i>	Gellérthegy	04.02. 1995
	<i>Ligustrum vulgare</i>	Tabán	27.01. 1995
	<i>Malus baccata</i>	Gellérthegy	08.05. 1991
	<i>Malus halliana</i>	Tabán	24.01. 1995
	<i>Malus × purpurea</i>	Vérmező	05.05. 1992
	<i>Phellodendron amurense</i>	Margitsziget	05.05. 1992
	<i>Populus alba</i>	Budafok	21.02. 1991
		Órméző	07.02. 1995
	<i>Populus × canescens</i>	Óbuda	12.05. 1995
	<i>Populus simonii</i>	Vérmező	25.08. 1998
		Sasad	25.01. 1991
	<i>Prunus cerasifera</i>	Tabán	23.09. 1991
		Gellérthegy	07.02. 1991
	<i>Prunus fruticosus</i>	Zugló	08.05. 1991
	<i>Prunus spinosa</i>	Vérmező	18.03. 1995
	<i>Pyrus betulifolia</i>	Gellérthegy	07.03. 1993
	<i>Pyrus pashia</i>	Gellérthegy	18.01. 1995
	<i>Pyrus pyraster</i>	Józsefváros	18.01. 1995
	<i>Rhus typhina</i>	Kőbánya	04.01. 1995
	<i>Syringa vulgaris</i>	Tabán	04.01. 1995
			12.02. 1991
fam. Glycyphagidae			
*<i>Hericia hericia</i> Robin, 1868			
	<i>Platanus × hispanica</i>	Józsefváros	16.05. 1991
Acari: Oribatida			
fam. Micreremidae			
<i>Micreremus brevipes</i> (Michael, 1888)			
	<i>Crataegus monogyna</i>	Gellérthegy	05.07. 1996
	<i>Daphne mezereum</i>	Törökvesz	26.01. 1995
	<i>Malus spectabilis</i>	Vérmező	03.04. 1992
	<i>Prunus spinosa</i>	Lanterna (Croatia)	07.03. 1993
	<i>Quercus ilex</i>		29.03. 1995
fam. Oribatulidae			
<i>Phauloppiella conformis</i> (Berlese, 1895)			
	<i>Juglans nigra</i>	Kőbánya	

Table 1 (cont.)

Mite species	Plant species	Sampling place	Sampling date
fam. Cymbaeremaeidae			
<i>Scapheremaeus reticulatus</i> (Berlese, 1910)			
	<i>Ilex aquifolium</i>	Városliget	04.02. 1995
	<i>Malus pumila</i> var. <i>niedzwetzkyana</i>	Gellérthegy	26.01. 1995
fam. Ceratozetidae			
<i>Trichoribates trimaculatus</i> (C. L. Koch, 1836)			
	<i>Jasminum beesianum</i>	Gellérthegy	27.01. 1995
fam. Scheloribatidae			
<i>Scheloribates latipes</i> (C. L. Koch, 1844)			
	<i>Jasminum beesianum</i>	Gellérthegy	27.01. 1995

Discussion

The corticolous fauna of species Acari has not been so far investigated extensively in Hungary.

The studies of Lebrun (1976), André (1986), also Momen and Lundqvist (1996) testify that the expected mite species living in a relatively small area can appear surprisingly rich.

The bark and bark crevices are the most important overwintering sites of foliage-dwelling mites. Some of them viz. tetranychid, tenuipalpid, phytoseiid, stigmeid, tydeid, winterschmidtiid species overwinter under the shield of dead coccids (Ripka, 1998a, b, Ripka and Kaźmierski, 1998a, b). There is a closer association between some members of family Tarsonemidae and coccids. *Tarsonemus nodosus*, *T. lobosus*, *T. karli*, *T. waitei* and *Dendroptus willmanni* also an other *Dendroptus* sp. quite often were found together with scale insects (Ripka et al., 1997). Fungivorous tarsonemid mites prefer this microhabitat. They feed on microscopic fungi.

Some tydeid species gave definite preference to cortex.

The predaceous members of corticolous cameroobiid, cheyletid, cunaxid, eupalpsellid, hemisarcopid, stigmeid and tydeid mite fauna cannot play a great role in regulating numbers of phytophagous insects and mites because of their low densities. But the knowledge on the predatory activities of these mites is sparse and mostly brief.

Acknowledgements

Cunaxoides kielczewskii (Acari: Cunaxidae) was identified by Robert L. Smiley, (Systematic Entomology Laboratory, Agricultural Research Service, US Department of Agriculture, Beltsville). Our thanks are due to Mr. R. L. Smiley for identifying this species. Special thanks are also expressed to Dr. Sándor Mahunka (Hungarian Natural History Museum, Budapest) and Dr. Wojciech Ł Magowski (A. Mickiewicz University, Poznań) for identifying of oribatid and tarsonemid mites, respectively.

The study was supported by Ministry of Agriculture and Regional Development, Hungary.

Literature

- André, H. M. (1986): Notes on the ecology of corticolous epiphyte dwellers. 4. Actinedida (especially Tydeidae) and Gamasida (especially Phytoseiidae). *Acarologia* 27, 107–115.
- André, H. M. (1987): Tydeidae (Acaria: Tydeidae) from Belgium II. The genera *Tydeus*, *Idiolorryia* and *Metatorryia*. *Acarologia* 28, 151–159.
- Balogh, J. and Mahunka, S. (1980): Mites XV. – Acari XV. Oribatida I. Fauna Hung., 139 (8), Akadémiai Kiadó, Budapest, 1–177.
- Bolland, H. R. (1983): A description of *Neophyllobius aesculi* n. sp. and its developmental stages (Acari: Camerobiidae). *Entomologische Berichten* 43, 42–47.
- Bolland, H. R. (1986): Review of the systematics of the family Camerobiidae (Acari, Raphignathoidea). I. The genera *Camerobia*, *Decaphyllobius*, *Tillandsobius* and *Tycherobius*. *Tijdschrift voor Entomologie* 129, 191–215.
- Bolland, H. R. (1991): Review of the systematics of the Camerobiidae, II. The genus *Neophyllobius* Berlese, 1886 (Acari: Raphignathoidea). *Genus* 2, 59–226.
- Cooreman, J. (1958): Description de *Tarsonemus bachmaieri* n. sp. (Taronemidae) et note sur *Eupalopsellus ölandicus* Sellnick (Eupalopsellidae). *Bulletin of Institut royal des Sciences naturelles de Belgique* 34, 1–8.
- Doberski, J. (1986): Population dynamics of corticolous mites of the genus *Daidalotarsonemus* (Acari: Tarsonomidae) on elm coppice. *Acarologia* 28, 31–36.
- Fain, A. (1982): Revision des genres *Thyreophagus* Rondani, 1874 et *Michaelopus* Fain et Johnston, 1974 (Acari, Acaridae) avec description de neuf espèces nouvelles. *Bull. Inst. r. Sci. nat. Belg.* 54, 1–47.
- Fain, A., Hurst, G. D. D., Tweddle, J. C., Lachlan, R. F., Majerus, M. E. N. and Britt, D. P. (1995): Description and observations of two new species of Hemisarcopidae from deutonymphs phoretic on Coccinellidae (Coleoptera) in Britain. *Internat. J. Acarol.* 21, 99–106.
- Fain, A. and Ripka, G. (1998): A new species of *Hemisarcopes* Lignières, 1893 (Acari: Hemisarcopidae) from ornamental trees in Hungary. *Internat. J. Acarol.* 24, 33–39.
- Gerson, U. (1973): The mites associated with armored scale insects. In: Daniel, M. and Rosicky, B. (eds): Proceedings of the 3rd International Congress of Acarology, Prague, 653–654.
- Gerson, U. (1985): Other predaceous mites and spiders. In: Helle, W. and Sabelis, M. W. (eds): Spider mites their biology, natural enemies and control. Elsevier, Amsterdam, Oxford, New York, Tokyo. 205–210.
- Kehl, C. and Weigmann, G. (1992): Die Hornmilbenzönosen (Acari, Oribatida) an Apfelbäumen im Stadtgebiet von Berlin als Bioindikatoren für die Luftqualität. *Zool. Beitr. N. F.* 34, 261–271.
- Keifer, H. H. (1952): Eriophyid Studies XIX. The Bulletin Department of Agriculture State of California. 41, 65–74.
- Keifer, H. H. (1975): Eriophyoidea Nalepa. In: Jeppson, L. R., Keifer, H. H. and Baker, E. W. (eds): Mites Injurious to Economic Plants. University of California Press, Berkeley, Los Angeles, London, 327–397.
- Komlovszky, I. Sz. (1980): A dendrofil atkák minőségi és mennyiségi viszonyai. Kandidáusi értekezés. (Qualitative and quantitative relations of dedrophil mites. PhD thesis). Szarvas–Budapest, 114 + xxx pp. (in Hungarian).
- Komlovszky, I. Sz. (1984): A fenyők (Coniferopsida) kártevő és ragadozó atka fajai. (Phytophagous and predatory mites on conifers [Coniferopsida]). *Növényvédelem* 20, 166–173. (in Hungarian).
- Lebrun, Ph. (1976): Effets écologiques de la pollution atmosphérique sur les populations et communautés de microarthropodes corticoles (Acariens, Collemboles et Ptérygotes). *Bull. Ecol.* 7, 417–430.
- Lucza, Z., Ripka, G. and Saly, K. R. (1996): Data to the Cheyletidae (Acari: Prostigmata) fauna of Hungary. *Folia Entomologica Hungarica* 57, 105–108.
- Meyer, M. K. P. (Smith) and Ueckermann, E. A. (1984): The family Eupalosellidae (Acari: Prostigmata), with descriptions of new species from South Africa. *Phytophylactica* 16, 121–142.
- Momen, F. M. (1987): The mite fauna of an unsprayed apple orchard in Ireland. *Angewandte Zoologie* 4, 417–431.

- Momen, F. M. (1988): New species of mites of the family Tydeidae (Acarina: Prostigmata) collected from unsprayed apple trees in Ireland. *Acarologia* 29, 355–360.
- Momen, F. and Lundqvist, L. (1996): Taxonomy of non-*Tydeus* genera of mite family Tydeidae (Acaria: Prostigmata) from moss, lichens and trees in southern Sweden. *Acarologia* 37, 281–297.
- Nachev, P. D. and Trenchev, G. (1987): The acarofauna associated with some Coccidae in Bulgaria. *Pochvoznanie, Agrokhimiya i Rastitelna Zashchita* 22, 81–86. (in Bulgarian).
- Ripka, G. (1998a): New data to the knowledge on the phytoseiid fauna in Hungary (Acaria: Mesostigmata). *Acta Phytopathologica et Entomologica Hungarica* 33, 395–405.
- Ripka, G. (1998b): New data to the knowledge on the tetranychid and tenuipalpid fauna in Hungary (Acaria: Prostigmata). *Acta Phytopathologica et Entomologica Hungarica* 33, 425–433.
- Ripka, G. and Kaźmierski, A. (1998a): New data to the knowledge on the tydeid fauna in Hungary (Acaria: Prostigmata). *Acta Phytopathologica et Entomologica Hungarica* 33, 407–418.
- Ripka, G. and Kaźmierski, A. (1998b): New data to the knowledge on the stigmaeid fauna in Hungary (Acaria: Prostigmata). *Acta Phytopathologica et Entomologica Hungarica* 33, 419–424.
- Ripka, G. and de Lillo, E. (1997): New data to the knowledge on the eriophyoid fauna in Hungary (Acaria: Eriophyoidea). *Folia Entomologica Hungarica* 58, 147–157.
- Ripka, G., Magowski, W. L. and Reider, K. (1997): Recent data on the knowledge of the fauna of tarsonemid mites (Acaria: Heterostigmata) on ornamental trees and shrubs. *Folia Entomologica Hungarica* 58, 159–168.
- Ripka, G., Saly, K. R. and Kozár, F. (1996): Újabb adatok a díszfa- és díszcserjefajok pajzstető- és lisztteske- (Homoptera: Coccoidea, Aleyrodoidea) faunájának ismeretéhez a fővárosban és környékén. (Recent data to the coccid and aleyrodid fauna [Homoptera: Coccoidea, Aleyrodoidea] on woody ornamentals in the Budapest area.) *Növényvédelem* 32, 7–17. (in Hungarian with English summary)
- Schmutterer, H. (1959): Schildläuse oder Coccoidea. In: Dahl, M. und Bischoff, H. (eds): *Die Tierwelt Deutschlands*. Gustav Fischer Verlag, Jena, 45, 260 pp.
- Smiley, R. L. (1992): The predatory mite fauna Cunaxidae (Acaria) of the world: with a new classification. Indira Publishing House, West Bloomfield, 356 pp.
- Smiley, R. L. and Knutson, L. (1983): Aspects of taxonomic research and services relative to mites as biological control agents. In: Hoy, M. A., Cunningham, G. L. and Knutson, L. (eds): *Biological Control of Pests by Mites*. USDA, 148–164.
- Suski, Z. W. (1972): Tarsonemid mites on apple trees in Poland. XI. Field observations on the distribution and significance of Tarsonemidae (Acarina, Heterostigmata) in apple orchards. *Zesz. Probl. Post. Nauk Roln.* 129, 139–157.
- Vacante, V. (1989): Gli Acari degli agrumi in Italia. III.: Gli Acarididi (Hemisarcopidae Oudemans, Acaridae Ewing and Nesbitt, Saprophlyidae Oudemans e Glycyphagidae Berlese). *Boll. Zool. agr. Bachic.* 2, 45–84.
- Vacante, V. and Gerson, U. (1987): Three species of *Eryngiopus* (Acaria: Stigmeidae) from Italy, with key to species and summary of habitats. *Redia* 70, 385–401.
- Weigmann, G. and Jung, E. (1992): Die Hornmilben (Acaria, Oribatida) an Strassenbäumen in Stadtzonen unterschiedlicher Luftbelastung in Berlin. *Zool. Beitr. N. F.* 34, 273–287.

