

HYPERPARASITISM OF STORKS
BY HYPOPI OF HYPODERIDAE
WITH DESCRIPTION OF A NEW SPECIES
OF THE GENUS **Neottialges**
(Acarina : Sarcoptiformes)

by A. FAIN*

INTRODUCTION

Hypopi belonging to the genus *Neottialges* have been found by Dr J.M. TUFTS, (Veterinary department, Ralston Purina City, St-Louis, U.S.A.), in enormous number, under the skin of three storks which died in the Zoo of St-Louis, Missouri, U.S.A.

The cause of the death of these birds was not clear. Postmortem examination did not reveal any significant gross lesions or any evidence of the cause of death other than heavy concentrations of this peculiar mite distributed throughout the subcutaneous tissues of all three birds. Concentrations of the mites were especially heavy on the fascia of the thighs. The birds apparently did show signs of leg weakness prior to death. One of the birds was in thin condition, but the other two appeared to be in fair flesh. Considerable water was present in the respiratory tract of one bird which apparently got down in the water prior to death. The wood stork is a native bird and the painted storks were imported from Europe about six months prior to death of the birds. (Dr J.M. TUFTS, *in litt.*)

These hypopi belong to two different species of Hypoderidae, from which one is new and is described here.

* Institut de Médecine Tropicale, Nationalestraat 155, 2000 Antwerpen.

FAMILY HYPODERIDAE MURRAY, 1877

Genus **Neottialges** FAIN, 1966

Subgenus **Caloenectes** FAIN, 1966

1. **Neottialges (Caloenectes) kutzeri** FAIN, 1967

This species has been described from *Ibis leucocephalus*. This bird died in the Zoo of Vienna, Austria.

About two thirds of the total number of hypopi found in the skin of the painted stork (*Ibis leucocephalus*), from the Zoo of St-Louis, U.S.A., belong to this species. The hypopi were very numerous. A small piece of skin of about 1 cm square, from the abdominal wall, contained several hundreds of hypopi belonging to this species.

2. **Neottialges (Caloenectes) ciconiarum** spec. nov.

This new species is distinguished from the other species in the subgenus by the following characters : setae *d* 2 long, setae *d* 3 very short, presence of a long sclerotized longitudinal band in front of the epimera III and fused posteriorly with these epimera.

HYPOPUS (fig. 1-5) : The holotype is 816 μ long and 330 μ wide. In 1 paratype (length x wide) : 840 μ x 355 μ . The sejugal furrow is present dorsally. Cuticle uniformly punctate except along the sejugal furrow and two postero-lateral longitudinal and curved lines (= ecdysis lines). The sclerotization of the cuticle is more marked in the regions of the posterior coxae, the propodosoma dorsally and the palposoma area. Epimera I fused in Y ; other epimera free. Genital suckers subequal distinctly divergent and separate by a strongly sclerotized area, a true genital sclerite however is absent. Anus very small situated 51 μ behind the genital suckers. Tarsi I and II long, 30 μ and 39 μ , respectively, ending in a slightly recurved spine, shorter (19 μ) than the respective tarsi. Tarsus III 69 μ long, ending in a slightly curved spine. Tarsus IV 51 μ long, with a strong and long barbed apical hair.

Chaetotaxy of idiosoma : setae *sc i*, *sc e*, *d* 1, *d* 2, *d* 5, *l* 1, *h*, *sh* very thin and respectively 60 μ , 200 μ , 300 μ , 200 μ , 180 μ ,

250-300 μ , 150 μ and 120 μ . Setae *d* 3, *d* 4, *l* 2, *l* 3, *l* 4, *l* 5 very thin and very short (less than 15 μ).

Chaetotaxy of legs : tarsus IV with 3 spines and 1 strong barbed apical hair, 300 μ long.




FIG. 1-2. —*Neottialges (Caloenectes) ciconiarum* sp.n. Hypopi in ventral and dorsal view.

Host and locality :

On a painted stork. *Ibis leucocephalus*, which died in the St Louis Zoo in August 1972 (Holotype and numerous paratypes).

The other storks (one painted stork and one wood stork *Mycrotergia americana* were parasitized by apparently the same hypopi but the latter have not been examined.

Types and paratypes in the U.S. National Museum, Washington. Paratypes in Institut royal des Sciences naturelles de Belgique and in the collection of the author.

Pathological action of these hypopi.

The hypopi have been found in enormous quantities in the abdominal walls and along the fascia of the thighs. It has not been




FIG. 3-5. — *Neottialges (Caloenectes) ciconiarum* sp.n. Hypopi. Fig. 1 : tarsi, tibia and genu of legs I ; fig. 2 : of leg III ; fig. 3 : of leg IV.

investigated in detail if they were present in other parts of the body.

It is well known that in some cases the hypopi of Hypoderidae may invade the connective tissue of any place of the body walls and of the viscera. We have found such hypopi previously in the following localisations, e.g. under the nasal mucosa, around the oesophagus, in the rachis of the feathers, in the air-sacs, on the surface of the lungs, in the lungs, in the gular pouch tissue, attach-

ed to the walls of the main blood vessels of the pleural cavity, between the pericardium and the heart muscles, etc. (FAIN and HYLAND, 1962, FAIN, 1967, FAIN and AMERSON, 1968).

It is not certain whether their presence in these localisations is always harmless. One may surmise that they could produce some toxic effect when they are extremely numerous as it was the case here. They also could produce local impairment when they invade the superficial layers of some vital organ such as the heart and the main blood vessels.

ACKNOWLEDGMENTS

I thank very much D^r J.M. TUFTS, Senior Veterinary Pathologist of the Ralston Purina Cy, St-Louis, Missouri, U.S.A. who sent me the hypopi and provided me with data concerning the occurrence of these parasites.

I also wish to express my gratitude to D^r C. YUNKER, Rocky Mountain Laboratory, who allowed the forwarding of this material to me.

BIBLIOGRAPHY

- FAIN A., 1967. — Les Hypopes parasites des tissus cellulaires des Oiseaux (Hypodectidae : Sarcoptiformes). *Bull. Inst. r. Sci. nat. Belg.*, 43 (4) : 1-139.
- FAIN A. and HYLAND K., 1962. — The Mites parasitic in the lungs of Birds. The variability of *Sternostoma tracheacolum* LAWRENCE, 1948, in domestic and wild birds. *Parasitology*, 52, (3-4) : 401-424.
- FAIN A. and AMERSON A. BINION jr., 1968. — Two new heteromorphic deutonymphs (Hypopi) (Acarina : Hypoderidae) from the Great Frigate-bird (*Fregata major*). *J. Med. Ent.*, 5 (3) : 320-324.
-